

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/234935632>

First Case of Chytridiomycosis in an Adult Specimen of a Native Anuran from Uruguay

Article in *Herpetological Review* · March 2011

CITATIONS

14

READS

177

4 authors, including:



Romina Ghirardi

Instituto Nacional de Limnología (INALI)

40 PUBLICATIONS 335 CITATIONS

[SEE PROFILE](#)



Raul Maneyro

Universidad de la República de Uruguay

112 PUBLICATIONS 2,113 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Comparative phylogeography of amphibians and reptiles of the Pampas [View project](#)



In Search of the Giant of the Pampas [View project](#)

—, M. D. McLEAN, AND S. BUCHANAN. 2009. Oviposition in northern clade *B. fowleri*: implications for conservation. *Appl. Herpetol.* 6:343–353.

WELDON, C., L. H. DU PREEZ, A. D. HYATT, R. MULLER, AND R. SPEARE. 2004. Origin of the amphibian chytrid fungus. *Emerg. Infect. Dis.* 10:2100–2105.

YOUNG, S., L. BERGER, AND R. SPEARE. 2007. Amphibian chytridiomycosis: strategies for captive management and conservation. *Int. Zoo Yb.* 41:85–95.

ZAR, J. H. 1999. *Biostatistical Analysis*. Prentice Hall, Englewood Cliffs, New Jersey. 663 pp.

Herpetological Review, 2011, 42(1), 65–66.
© 2011 by Society for the Study of Amphibians and Reptiles

First Case of Chytridiomycosis in an Adult Specimen of a Native Anuran from Uruguay

In Uruguay, the amphibian chytrid fungus *Batrachochytrium dendrobatidis* (*Bd*) was first detected by Mazzoni et al. (2003). They reported chytridiomycosis in a captive population of *Lithobates catesbeianus* (Anura, Ranidae) which suffered a massive death of metamorphs at a commercial farm. Later, Garner et al. (2006) also confirmed the presence of *Bd* at the same farm and in a new one. Subsequently, Borteiro et al. (2009) found *Bd* by histological analysis in the oral epithelium of native wild species larvae: *Odontophrynus maisuma* (Anura, Cycloramphidae), *Physalaemus henselii* (Anura, Leiuperidae), *Hypsiboas pulchellus* and *Scinax squalirostris* (Anura, Hylidae).

We sampled native, wild amphibians for *Bd* in June 2009 at the Cerro Verde Protected Area, Rocha Department, Uruguay (33.945833°S, 53.508611°W) (Fig. 1). This area is characterized by humid subtropical climate (average temperature 16°C, annual rainfall 950 mm) and an ocean coastline with dunes (20–30 m elevation) and plains (Alonso and Bassagoda 2003). During nighttime sampling, *Pleurodema bibroni* were heard calling at a temporary pond of a coastal meadow. One frog was collected as a voucher specimen of the survey and another individual was collected under suspicion of being infected by *Bd* because it possessed some lesions on the skin (two ulcerations: on right side and dorsum, as described by Berger et al. 2005). Both specimens were taken to the laboratory and the injured frog died a few days later. Both were deposited at Vertebrate Zoology Collection, Facultad de Ciencias, Universidad de la República (ZVC-B 19776, ZVC-B 19775).

To assess the presence of *Bd* in both frogs, skin of the underside of the legs, interdigital membrane of feet, and drink patch was comprehensively swabbed (10 times ea.). Each swab was broken off into an empty Eppendorf tube, and then analyzed




FIG. 1. Location of the Cerro Verde Protected Area, Rocha Department, Uruguay, where *Batrachochytrium dendrobatidis* was detected on *Pleurodema bibroni* (star).

CECILIA BARDIER
Instituto de Ecología y Ciencias Ambientales, Facultad de Ciencias (UdelaR) Iguá 4225 (11400), Montevideo, Uruguay
e-mail: ceciliabardier@gmail.com

ROMINA GHIRARDI
Instituto Nacional de Limnología (INALI; CONICET-UNL)
Ciudad Universitaria, Paraje El Pozo (3000)
Santa Fe, Argentina
and
Instituto de Botánica Spegazzini (FCNyM-UNLP)
Calle 53 N° 477 (1900), La Plata, Buenos Aires, Argentina
e-mail: rghirardi@inali.unl.edu.ar

MICHAEL LEVY
Department of Population Health and Pathobiology
North Carolina State University, College of Veterinary Medicine
Raleigh, North Carolina 27606, USA
e-mail: mglevy@ncsu.edu

RAÚL MANEYRO
Instituto de Ecología y Ciencias Ambientales, Facultad de Ciencias (UdelaR) Iguá 4225 (11400), Montevideo, Uruguay
e-mail: rmaneyro@fcien.edu.uy




FIG. 2. Skin section of *Pleurodema bibroni* (ZVC-B 19775), collected from Cerro Verde Protected Area, Rocha Department, Uruguay, showing empty *Batrachochytrium dendrobatidis* zoosporangium (EZ) and zoosporangium containing zoospores (Z), one with a discharge tube (DT). Scale bar: 20 μ m.

for the presence of *Bd* DNA using qPCR (Boyle et al. 2004). *Bd* infection was also analyzed histologically. Abdominal and hind limb ventral skin patches (~5 × 10 mm) and two toe tips were excised from the anurans and routinely processed for histological examination following Berger et al. (1999).

Both analyses (qPCR and histology) were *Bd*-positive only for the apparently healthy specimen. Zoospores and zoosporangia at different stages were found at multiple sections of toe tips and hind limb of the infected individual, and some degree of hyperkeratosis at the sections from hind limbs of the specimen was found (Fig. 2).

This is the first record of *Bd* in an adult native wild-caught amphibian in Uruguay, and the first report of *Bd* infection for *Pleurodema bibroni*. This frog is considered Near Threatened by the IUCN Red List (IUCN 2006) and is Threatened at a country level for Uruguay (Maneyro and Langone 2001; Canavero et al. 2010). The pathogen was detected 100 km northeast of the nearest record in Uruguay (Borteiro et al. 2009) inside a Protected Area, which is both alarming and suggests that measures for control and prevention of human-mediated *Bd* dispersal warrant consideration. This is particularly important because populations of other native species in the area, such as *Melanophryniscus montevidensis* (Núñez et al. 2004), may be at risk (IUCN RED LIST: Vulnerable; IUCN 2006).

Acknowledgments.—We thank M. Bessonart and F. Féola from the Facultad de Ciencias for their support for histological analysis at the Estación Experimental de Investigaciones Marinas y Acuicultura, K. Gore and M. Perdue for molecular analysis, F. Gallego for graphical assistance, and I. Berro and A. García for invaluable help during field surveys.

Resumen.—En Uruguay la infección causada por *Batrachochytrium dendrobatidis* (*Bd*) había sido detectada en *Lithobates catesbeianus* de criaderos comerciales y en larvas silvestres de especies autóctonas, pero no había sido confirmada aún en ejemplares adultos de ninguna especie nativa. En junio de 2009 se colectaron dos machos adultos silvestres de *Pleurodema bibroni* en el Cerro Verde (33.935°S, 53.5070556°W, Rocha, Uruguay), se les tomaron muestras para análisis de ADN (qPCR) para detectar *Bd*. Uno de ellos resultó positivo y el análisis histológico de la piel del ejemplar también confirmó la infección. Por tanto se diagnostica por primera vez la infección por *Bd* en adultos silvestres de esta especie en Uruguay, que además está considerada como *Amenazada* a nivel nacional y *Near Threatened* para la IUCN.

LITERATURE CITED

- ALONSO, E., AND M. BASSAGODA. 2003. Relevamiento de la flora y comunidades vegetales del Cerro Verde, Rocha, Uruguay. *Comunicaciones Botánicas. Museo Nacional de Historia Natural y Antropología* 127:1–20.
- BERGER, L., A. D. HYATT, R. SPEARE, J. E. LONGCORE. 2005. Life cycle stages of the amphibian chytrid *Batrachochytrium dendrobatidis*. *Dis. Aquat. Org.* 68(1):51–63.
- , R. SPEARE, AND A. KENT. 1999. Diagnosis of chytridiomycosis in amphibians by histologic examination. Available at: <http://www.jcu.edu.au/school/phtm/PHTM/frogs/histo/chhisto.htm>.
- BORTEIRO, C., J. CRUZ, F. KOLENC, AND A. ARAMBURU. 2009. Chytridiomycosis in frogs from Uruguay. *Dis. Aquat. Org.* 84:159–162.
- BOYLE, D. G. D. B. BOYLE, V. OLSEN, J. A. T. MORGAN, AND A. D. HYATT. 2004. Rapid quantitative detection of chytridiomycosis (*Batrachochytrium dendrobatidis*) in amphibian samples using real-time Taqman PCR assay. *Dis. Aquat. Org.* 60(2):141–148.
- CANAVERO, A., S. CARREIRA, J. A. LANGONE, F. ACHAVAL, C. BORTEIRO, A. CAMARGO, I. DA ROSA, A. ESTRADES, A. FALLABRINO, F. KOLENC, M. M. LÓPEZ-MENDILAHARSU, R. MANEYRO, M. MENEGHEL, D. NUÑEZ, C. M. PRIGIONI, AND L. ZIEGLER. 2010. Red list of the amphibians and reptiles of Uruguay. *Iheringia (Ser. Zool.)* 100:5–12.
- GARNER, T. W., M. W. PERKINS, P. GOVINDARAJULU, D. SEGIE, S. WALKER, A. A. CUNNINGHAM, AND M. C. FISHER. 2006. The emerging amphibian pathogen *Batrachochytrium dendrobatidis* globally infects introduced populations of North American bullfrog, *Rana catesbeiana*. *Biol. Lett.* 2:455–459.
- IUCN, CONSERVATION INTERNATIONAL AND NATURESERVE. 2006. Global Amphibian Assessment. Available at: www.globalamphibians.org. Accessed 28 July 2009.
- MANEYRO, R., AND J. A. LANGONE. 2001. Categorización de los anfibios del Uruguay. *Cuad. Herpetol.* 15(2):107–118.
- MAZZONI, R., A. A. CUNNINGHAM, P. DASZAK, A. APOLO, E. PERDOMO, AND G. SPERANZA. 2003. Emerging pathogen of amphibians in frogs (*Rana catesbeiana*) farmed for international trade. *Emerg. Infect. Dis.* 9:995–998.
- NUÑEZ, D., R. MANEYRO, J. A. LANGONE, AND R. O. DE SÁ. 2004. Distribución geográfica de la fauna de anfibios del Uruguay. *Smithson. Herpetol. Infor. Serv.* 134:1–36.