

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/26247582>

# Chytridiomycosis in frogs from Uruguay

Article in *Diseases of Aquatic Organisms* · May 2009

DOI: 10.3354/dao02035 · Source: PubMed

---

CITATIONS

20

---

READS

187

4 authors, including:



**Claudio Borteiro**

Museo Nacional de Historia Natural de Montevideo

76 PUBLICATIONS 945 CITATIONS

[SEE PROFILE](#)



**Juan Carlos Cruz**

Universidad Politécnica Metropolitana de Hidalgo

19 PUBLICATIONS 131 CITATIONS

[SEE PROFILE](#)



**Francisco Kolenc**

Museo Nacional de Historia Natural, Montevideo, Uruguay

73 PUBLICATIONS 910 CITATIONS

[SEE PROFILE](#)

NOTE

## Chytridiomycosis in frogs from Uruguay

Claudio Borteiro<sup>1,\*</sup>, Juan Carlos Cruz<sup>2</sup>, Francisco Kolenc<sup>3</sup>, Andrea Aramburu<sup>4</sup>

<sup>1</sup>Río de Janeiro 4058, CP 12800, Montevideo, Uruguay

<sup>2</sup>Departamento de Patología, Facultad de Veterinaria, Lasplacas 1550, Montevideo, Uruguay

<sup>3</sup>Universidad Católica del Uruguay, Dr. Javier Barrios Amorín 1578, Montevideo, Uruguay

<sup>4</sup>Calle 43 y Calle O, Parque del Plata, Canelones, Uruguay

**ABSTRACT:** Amphibian chytridiomycosis caused by *Batrachochytrium dendrobatidis* is reported in Uruguayan native amphibians for the first time. Histological evidence of infection was observed in tadpoles of *Hypsiboas pulchellus*, *Odontophrynus maisuma*, *Physalaemus henselii*, and *Scinax squalirostris*. The effects of chytridiomycosis on these species are still unknown. However, the disease is of potential concern for the conservation of the apparently declining species *P. henselii* and also for *O. maisuma*, given its restricted distribution in habitats which are being increasingly disturbed.

**KEY WORDS:** *Batrachochytrium dendrobatidis* · Amphibian disease · Amphibian conservation · Uruguay

Resale or republication not permitted without written consent of the publisher

### INTRODUCTION

Many amphibian species have disappeared or are severely threatened because of multiple factors such as habitat loss, pollution, introduction of non-native species, climate change and infectious diseases (Stuart et al. 2004, Pounds et al. 2006). Studies that focused on pathogens as a cause of amphibian declines greatly increased after the discovery of amphibian chytridiomycosis, a fungal skin disease caused by the chytrid fungus *Batrachochytrium dendrobatidis* (see Daszak et al. 2007). Chytridiomycosis can be highly pathogenic to amphibian non-larval stages and is linked with mass mortalities and population declines (Berger et al. 1998, Daszak et al. 1999, 2003). The involvement of chytridiomycosis in the decline and extinction of Neotropical amphibians was first documented in Central America and northern South America, where it strongly impacted several amphibian communities (Berger et al. 1998, Lips et al. 2003, 2006). In recent years, *B. dendrobatidis* has been reported to be widespread farther south, principally along the Atlantic coast of South

America, where disease-linked declines are suspected (Barrionuevo & Mangione 2006, Carnaval et al. 2006, Toledo et al. 2006). Chytridiomycosis is likely to occur in Uruguayan native amphibians because climatic conditions are highly suitable for the pathogen to be established in the region (Ron 2005). Furthermore, the disease has already been reported from this country in farmed North American bullfrogs *Lithobates catesbeianus* (Mazzoni et al. 2003).

The amphibian chytrid is associated with keratinized tissues, thus detection is made on samples of adult amphibian skin and oral epithelium surrounding keratinized mouthparts of tadpoles (Berger et al. 1999). Infection at the larval life stage essentially consists of variable amounts of keratin loss that may be non-lethal (Knapp & Morgan 2006, Symonds et al. 2007). When examining tadpoles of amphibians from northern Uruguay, we noticed the presence of chytrid-like deformities in the oral structures of tadpoles of the leiupeid frog *Physalaemus henselii*. This finding prompted us to screen for amphibian chytridiomycosis in this and other Uruguayan native amphibians.

\*Email: borteiro@gmail.com

Herein, we report new cases of the disease in this country, expanding its geographic and host range.

## MATERIALS AND METHODS

Twenty-seven tadpoles collected in Uruguay between 2001 and 2007 that exhibited deformities in keratinized oral structures under a stereoscopic microscope were screened for the presence of *Batrachochytrium dendrobatidis*. Vouchers were deposited in the Batrachians Collection of the Vertebrate Zoology Department (ZVCB), Faculty of Sciences, Montevideo, Uruguay. Specimen details are listed in Table 1. Tadpoles were fixed *in situ* with formalin 10%. The cephalic region of each specimen was dissected and routinely processed for histological examination; they were embedded in paraffin, sectioned at 4 to 5  $\mu\text{m}$  and stained with hematoxylin and eosin. Diagnosis followed the work of Berger et al. (1999).

## RESULTS

We detected chytrid on 18 out of 31 tadpoles: all screened specimens of *Hypsiboas pulchellus* (n = 6), *Odontophrynus maisuma* (4), *Scinax squalirostris* (3) and those of *Physalaemus henselii* from Pueblo Madera (5). Most relevant oral deformities of tadpoles were partial or total loss of keratinization of jaw sheaths, with erosion of their cutting edge and absence of serrations, and also stunted and missing teeth. The loss of labial teeth was sometimes so extensive that it resulted in wide gaps within tooth rows, or even the lack of entire tooth rows, especially anterior ones. Infected tadpoles were not emaciated, their general




Fig. 1. Section of the oral epithelium (oe) in a *Hypsiboas pulchellus* tadpole. Empty zoosporangia (z) of *Batrachochytrium dendrobatidis* are evident; note the septum (s) and discharge papillae (dp). Scale bar = 20  $\mu\text{m}$

aspect made them indistinguishable from non-infected tadpoles, and no other concurrent diseases or external abnormalities were observed. The diagnosis was based on the observation of various developmental stages of *Batrachochytrium dendrobatidis*, mainly mature zoosporangia with zoospores and empty zoosporangia (which were the most common stage), sometimes exhibiting discharge papillae and septae (Fig. 1).

## DISCUSSION

This is the first report of amphibian chytridiomycosis in wild amphibians from Uruguay, and also for the species *Hypsiboas pulchellus*, *Odontophrynus maisuma*, *Physalaemus henselii* and *Scinax squalirostris*. *Batrachochytrium dendrobatidis* was detected in tadpoles

Table 1. Details of specimens examined in the present study. ZVCB: Batrachians Collection of the Vertebrate Zoology Department, Faculty of Sciences, Montevideo, Uruguay. m.a.s.l.: meters above sea level; -: no voucher specimen deposited

Species	n	Locality	Lat./Long.	Altitude (m a.s.l.)	Collection date	ZVCB voucher no.
<i>Hypsiboas pulchellus</i>	4	Road from La Paloma to Laguna de Rocha, Rocha	34° 38' S, 54° 12' W	54	7 Aug 2006	16143
<i>H. pulchellus</i>	2	Delta del Tigre, San José	34° 46' S, 56° 21' W	5	2 Sep 2005	16144
<i>Melanophryniscus montevidensis</i>	3	Laguna de Rocha, Rocha	34° 39' S, 54° 13' W	0	10 Mar 2007	-
<i>M. sanmartini</i>	2	Sierra de las Ánimas, Maldonado	34° 44' S, 55° 19' W	450	15 Aug 2004	16141
<i>Odontophrynus maisuma</i>	4	Laguna de Rocha, Rocha	34° 39' S, 54° 13' W	0	7 Aug 2006	16142
<i>Physalaemus biligonigerus</i>	1	Quebrada de los Cuervos, Treinta y Tres	32° 57' S, 54° 27' W	185	4 Mar 2007	15192
<i>P. henselii</i>	5	Pueblo Madera, Rivera	30° 58' S, 55° 34' W	250	9 Oct 2001	16146
<i>P. henselii</i>	3	Barra de Valizas, Rocha	34° 20' S, 53° 48' W	2	10 Mar 2007	16149
<i>Scinax squalirostris</i>	3	Laguna de Rocha, Rocha	34° 39' S, 54° 13' W	0	7 Aug 2006	16145



Fig. 2. Geographic distribution of *Batrachochytrium dendrobatidis* in Uruguay. (★) New records in this study: (1) Pueblo Madera, (2) Delta del Tigre, (3) Road to Laguna de Rocha and Laguna de Rocha. (●) Previous reports in farmed bullfrogs *Lithobates catesbeianus*: (4) Libertad (Mazzoni et al. 2003), (5) Empalme Olmos (Garner et al. 2006)

collected during austral winter and early spring, from August to October. Tadpoles of *O. maisuma* and *S. squalirostris* positive for chytridiomycosis were collected in the same pond where tadpoles of *Melanophryniscus montevidensis* were apparently free of infection in late summer (March). This may be due to the small sample of *M. montevidensis* we studied or to the sensitivity of the histological diagnosis, but could also be explained by specific differential susceptibility of tadpoles (Blaustein et al. 2005) and/or seasonality (Kriger & Hero 2007). Mouthpart deformities observed in infected tadpoles are similar to those reported for amphibian chytridiomycosis elsewhere (Knapp & Morgan 2006). Mouthpart deformities previously reported in tadpoles of *P. henselii* (Kolenc et al. 2006) were due to chytrid infection, as confirmed in the present study.

Most reports about chytrid-driven declines in South America conclude that amphibian populations at high elevations are more threatened than those inhabiting lowlands (Berger et al. 1998, La Marca et al. 2005, Barrionuevo & Mangione 2006, Carnaval et al. 2006). However, reports of infection from coastal Uruguay (present study) and also from lowlands in Argentina (Herrera et al. 2005) indicate a wide altitudinal distribution of *Batrachochytrium dendrobatidis* in the region (see Table 1, Fig. 2). The presence of chytrid fungus in the population of *Physalaemus henselii* from northern

Uruguay is noteworthy as this species has apparently declined over much of its range in this country (F. Kolenc & C. Borteiro unpubl. data). In addition, the positive infection of *O. maisuma* is of special concern as this species has a narrow geographic distribution (Rosset 2008), mainly restricted to coastal habitats which are subject to disturbance due to rapid urbanization. During our surveys, adults of infected species studied herein were common at those sites where we detected *B. dendrobatidis* and dead adult amphibians were not found at any locality (F. Kolenc & C. Borteiro pers. obs.). As opposed to adult amphibians, the pathogenic effects of *B. dendrobatidis* on amphibian larval communities and in turn on their entire ecosystems are still poorly understood (Parris & Baud 2004, Ranvestel et al. 2004). We do not know the extent to which chytridiomycosis has impacted Uruguayan native amphibians, especially in coastal areas where amphibian declines caused by factors other than urbanization and habitat loss were recently reported (Kolenc et al. 2009).

**Acknowledgements.** We are grateful to M. Tedros, J. C. Borteiro, M. Pallas, G. Duarte and J. Valbuena for their hospitality and aid in surveys, P. Symonds for her suggestions, S. Ron, D. Martí and D. Baldo shared bibliography, and G. Duarte helped with image processing.

#### LITERATURE CITED

- Barrionuevo S, Mangione S (2006) Chytridiomycosis in two species of *Telmatobius* (Anura: Leptodactylidae) from Argentina. *Dis Aquat Org* 73:171–174
- Berger L, Speare R, Daszak P, Green DE and others (1998) Chytridiomycosis causes amphibian mortality associated with population declines in the rain forests of Australia and Central America. *Proc Natl Acad Sci USA* 95: 9031–9036
- Berger L, Speare R, Kent A (1999) Diagnosis of chytridiomycosis in amphibians by histologic examination. *Zoos Print J* 15:184–190
- Blaustein AR, Romansic JM, Scheessele EA, Han BA, Pessier AP, Longcore JE (2005) Interspecific variation in susceptibility of frog tadpoles of the pathogenic fungus *Batrachochytrium dendrobatidis*. *Conserv Biol* 19:1460–1468
- Carnaval ACOQ, Puschendorf R, Peixoto OL, Verdade VK, Trefaut Rodrigues M (2006) Amphibian chytrid fungus broadly distributed in the Brazilian Atlantic Rain Forest. *EcoHealth* 3:41–48
- Daszak P, Berger L, Cunningham AA, Hyatt AD, Green DE, Speare R (1999) Emerging infectious diseases and amphibian population declines. *Emerg Infect Dis* 5: 735–748
- Daszak P, Cunningham AA, Hyatt AD (2003) Infectious disease and amphibian population decline. *Divers Distrib* 9: 141–150
- Daszak P, Lips K, Alford R, Carey C and others (2007) Infectious diseases. In: Gascon C, Collins JP, Moore RD, Church DR, McKay JE, Mendelson JR III (eds) *Amphibian Conservation Action Plan*. IUCN/SSC, Gland, p 21–25
- Garner TWJ, Perkins MW, Govindarajulu P, Seglie D, Walker

- S, Cunningham AA, Fisher MC (2006) The emerging amphibian pathogen *Batrachochytrium dendrobatidis* globally infects introduced populations of North American bullfrog, *Rana catesbeiana*. *Biol Lett* 2:455–459
- Herrera RA, Steciow MM, Natale GS (2005) Chytrid fungus parasitizing the wild amphibian *Leptodactylus ocellatus* (Anura: Leptodactylidae) in Argentina. *Dis Aquat Org* 64: 247–252
- Knapp RA, Morgan JAT (2006) Tadpole mouthpart depigmentation as an accurate indicator of chytridiomycosis, an emerging disease of amphibians. *Copeia* 2006:188–197
- Kolenc F, Borteiro C, Tedros M, Núñez D, Maneyro R (2006) The tadpole of *Physalaemus henselii* (Peters) (Anura: Leiuperidae). *Zootaxa* 136:41–50
- Kolenc F, Borteiro C, Baldo D, Ferraro D, Prigioni C (2009) The tadpoles and advertisement calls of *Pleurodema bibroni* Tschudi and *Pleurodema kriegi* (Müller), with notes on their geographic distribution and conservation status (Amphibia, Anura, Leiuperidae). *Zootaxa* 1969:1–35
- Kruger KM, Hero JM (2007) Large-scale seasonal variation in the prevalence and severity of chytridiomycosis. *J Zool (Lond)* 271:352–359
- La Marca E, Lips KR, Lötters S, Puschendorf R and others (2005) Catastrophic population declines and extinctions in neotropical harlequin frogs (Bufonidae: *Atelopus*). *Biotropica* 37:190–201
- Lips KR, Green DR, Pappendick R (2003) Chytridiomycosis in wild frogs from southern Costa Rica. *J Herpetol* 37: 215–218
- Lips KR, Brem F, Brenes R, Reeve JD and others (2006) Emerging infectious disease and the loss of biodiversity in a Neotropical amphibian community. *Proc Natl Acad Sci USA* 103:3165–3170
- Mazzoni R, Cunningham AA, Daszak P, Apolo A, Perdomo E, Speranza G (2003) Emerging pathogen of wild amphibians in frogs (*Rana catesbeiana*) farmed for international trade. *Emerg Infect Dis* 9:995–998
- Parris MJ, Baud DR (2004) Interactive effect of a heavy metal and chytridiomycosis on Gray tree Frog larvae (*Hyla crysocelis*). *Copeia* 2004:344–350
- Pounds JA, Bustamante MR, Coloma LA, Consuegra JA and others (2006) Widespread amphibian extinctions from epidemic disease driven by global warming. *Nature* 439: 161–167
- Ranvestel AW, Lips KR, Pringle CM, Whiles MR, Bixby RJ (2004) Neotropical tadpoles influence stream benthos: evidence for the ecological consequences of decline in amphibian populations. *Freshw Biol* 49:274–285
- Ron S (2005) Predicting the distribution of the amphibian pathogen *Batrachochytrium dendrobatidis* in the New World. *Biotropica* 37:209–221
- Rosset SD (2008) New species of *Odontophrynus* Reinhardt and Lütken 1862 (Anura: Neobatrachia) from Brazil and Uruguay. *J Herpetol* 42:134–144
- Stuart SN, Chanson JS, Cox NA, Young BE, Rodrigues ASL, Fischman DL, Waller RW (2004) Status and trends of amphibian declines and extinctions worldwide. *Science* 306:1783–1786
- Symonds EP, Hines HB, Bird PS, Morton JM, Mills PC (2007) Surveillance for *Batrachochytrium dendrobatidis* using *Myxophyes* (Anura: Myobatrachidae) larvae. *J Wildl Dis* 43:48–60
- Toledo LF, Britto FB, Araújo OGS, Liasson LMO, Haddad CFB (2006) The occurrence of *Batrachochytrium dendrobatidis* in Brazil and the inclusion of 17 new cases of infection. *S Am J Herpetol* 1:185–191

*Editorial responsibility: David Marcogliese, Montreal, Quebec, Canada*

*Submitted: June 12, 2008; Accepted: January 14, 2009  
Proofs received from author(s): March 30, 2009*